

# SYMPLECTIC GROUPS ARE $N$ -DETERMINED 2-COMPACT GROUPS

ALEŠ VAVPETIČ AND ANTONIO VIRUEL

**ABSTRACT.** We show that for  $n \geq 3$  the symplectic group  $Sp(n)$  is as a 2-compact group determined up to isomorphism by the isomorphism type of its maximal torus normalizer. This allows us to determine the integral homotopy type of  $Sp(n)$  among connected finite loop spaces with maximal torus.

## 1. INTRODUCTION

The advent of  $p$ -compact groups in the celebrated work of Dwyer and Wilkerson [9] is the culmination of a research program that can be traced back to the work of Hopf and Serre on  $H$ -spaces and loop spaces, and fits within the philosophy of Hilbert's Fifth Problem: which are the non differential (here homotopy theoretical) properties that characterize compact Lie groups?

A  $p$ -compact group is a loop space  $(X, BX, e)$ , i.e.  $e : X \simeq \Omega(BX)$  for a pointed space  $BX$ , such that  $H^*(X; \mathbb{F}_p)$  is finite and  $BX$  is  $p$ -complete in the sense of Bousfield and Kan [4]. As expected, examples of  $p$ -compact groups are given by  $p$ -completion of compact Lie groups  $G$  for which  $\pi_0 G$  is a  $p$ -group, since  $G_p^\wedge$  is homotopy equivalent to  $\Omega(BG_p^\wedge)$ . In this way a  $p$ -compact torus  $T$  of rank  $n$  is the  $p$ -completion of an ordinary torus, hence  $BT$  is the Eilenberg-MacLane space  $K((\mathbb{Z}_p)^{\oplus n}, 2)$ . Further examples are given by the realization of polynomial algebras, i.e. loop spaces  $\Omega BX$ , where  $BX$  is  $p$ -complete and has polynomial mod- $p$  cohomology ([1], [5], [11], [27], [32], [37]). The importance of  $p$ -compact groups consists of a dictionary (reviewed in Section 2) that translates much of the rich internal algebraic structure of compact Lie groups to the homotopy theoretical setting of  $p$ -compact groups, so the challenge is then to give homotopy theoretical proofs of classical algebraic Lie group theory results.

One of those challenges quoted above is the following:  $p$ -compact groups admit maximal tori, Weyl groups and maximal torus normalizers in a way that extends the classical concepts in Lie group theory [9, 8.13 & 9.5], so can we “reproof” the Lie group theoretical Curtis-Wiederhold-Williams’ theorem [6] in the setting of  $p$ -compact groups? Recall that Curtis-Wiederhold-Williams’ theorem states that two compact connected Lie groups are isomorphic if and only if their maximal torus normalizers are isomorphic, hence we are led to the following conjecture [7, Conjecture 5.3]

---

*Date:* February 1, 2008.

The first author is partially supported by the Ministry for Education, Science and Sport of Republic of Slovenia Research Program no. 101-509. The second author is partially supported by the DGES-FEDER grant BFM2001-1825, and Junta de Andalucía Grant FQM-0213.

**Conjecture 1.1.** *Let  $X$  be a connected  $p$ -compact group with maximal torus  $T_X$ . Then  $X$  is determined up to equivalence by the loop space  $NT_X$ .*

We shall say that a  $p$ -compact group  $X$  is  $N$ -determined if  $X$  verifies Conjecture 1.1 even if the hypothesis “connected” is dropped, i.e.  $X$  is  $N$ -determined if every  $p$ -compact group  $Y$ , with the normalizer of a maximal torus isomorphic to that of  $X$ , is isomorphic to  $X$ .

Given an odd prime  $p > 2$ ,  $p$ -compact groups are known to be  $N$ -determined [2], what leads to the classification of  $p$ -compact groups for  $p$  odd. But the situation is quite different at  $p = 2$ : there exist 2-compact groups which are not  $N$ -determined. For example  $O(n)_2^\wedge$  and  $SO(n+1)_2^\wedge$  are non isomorphic 2-compact groups that have isomorphic maximal torus normalizers. So at  $p = 2$  we cannot drop the hypothesis “connected” in Conjecture 1.1.

We say that a 2-compact group  $X$  is weakly  $N$ -determined if every 2-compact group  $Y$ , for which there exists a homotopy equivalence  $BN_X \simeq BN_Y$  between the maximal torus normalizers of  $X$  and  $Y$ , inducing an isomorphism  $\pi_0 X \cong \pi_0 Y$ , is isomorphic to  $X$ . From the definitions it follows that an  $N$ -determined 2-compact group is also weakly  $N$ -determined.

It has been shown that the 2-compact groups  $O(n)_2^\wedge$ ,  $SO(2n+1)_2^\wedge$  and  $Spin(2n+1)_2^\wedge$  [25] are weakly  $N$ -determined 2-compact groups (and they are not  $N$ -determined), and that  $U(n)_2^\wedge$  for  $n \neq 2$  [23],  $(G_2)_2^\wedge$  [34],  $(F_4)_2^\wedge$  [33], and  $DI(4)$  [26] are  $N$ -determined ( $U(2)_2^\wedge$  is only weakly  $N$ -determined, because the normalizer  $N$  of a maximal torus of  $U(2)_2^\wedge$  is also a 2-compact group but  $N$  is not isomorphic to  $U(2)_2^\wedge$ ). In this paper we prove that the symplectic groups  $Sp(n)_2^\wedge$  are  $N$ -determined 2-compact groups for  $n \geq 3$ .

**Theorem 1.2.** *Let  $n \geq 3$  and let  $X$  be a 2-compact group with the maximal torus normalizer  $f_N: N \longrightarrow X$  isomorphic to that of  $Sp(n)_2^\wedge$ . Then  $X$  and  $Sp(n)_2^\wedge$  are isomorphic 2-compact groups.*

*Proof.* First we prove that  $X$  is connected in Section 3 (Proposition 3.1). In Section 4 we show that mod-2 cohomology of  $BX$  is isomorphic to that of  $BSp(n)$  as algebras over the Steenrod algebra, which implies that the Quillen categories associated to  $X$  and  $Sp(n)$  are isomorphic. In section 5 we describe the 2-stubborn decomposition of the group  $Sp(n)$ , which allows us to define a map from  $BSp(n)_2^\wedge$  to  $BX$  that happens to be an equivalence. This is done in Section 6.  $\square$

Notice that the hypothesis  $n \geq 3$  is necessary as  $Sp(1)_2^\wedge = SU(2)_2^\wedge$  and  $Sp(2)_2^\wedge = Spin(5)_2^\wedge$  are only weakly  $N$ -determined 2-compact groups.

The combination of the results in [2] and Theorem 1.2 shows that if  $G$  is a connected compact Lie group, then  $BG$  is in the adic genus of  $BSp(n)$  if and only if  $G = Sp(n)$ , what in view of [30] characterizes the integral homotopy type of  $BSp(n)$  as a loop space. Thus our final result is

**Theorem 1.3.** *Let  $L$  be a connected finite loop space with a maximal torus normalizer isomorphic to that of  $Sp(n)$ . Then  $BL$  is homotopy equivalent to  $BSp(n)$ .*

**Notation.** Here all spaces are assumed to have the homotopy type of a CW-complex. Completion means Bousfield-Kan completion [4]. For a given space  $X$ , we write  $H^*X$  for the mod-2 cohomology  $H^*(X; \mathbb{F}_2)$ . For a prime  $p$ , we write  $X_p^\wedge$  for the Bousfield-Kan  $p$ -completion ( $(\mathbb{Z}_p)_\infty$ -completion in the terminology of Bousfield and Kan) of the space  $X$ . We assume that the reader is familiar with Lannes' theory [17].

## 2. THE DICTIONARY

As announced in the introduction, this section is devoted to a brief review of the dictionary translating constructions and arguments from the algebraic theory of groups to the homotopical setting of  $p$ -compact groups. The aim of the minimalist style of this section is to ease the search of concepts by the reader who will find a more detailed exposition in the original [8], or the reviews [7], [21] and [28] if needed.

Along this section  $X$  and  $Y$  will denote  $p$ -compact groups whose classifying spaces are  $BX$  and  $BY$  respectively. By  $T$  we shall denote a  $p$ -compact torus, i.e.  $BT \simeq K(\mathbb{Z}_p^n, 2)$  where  $n$  is the rank of  $T$ . Finally, we define:

- **Homomorphisms** [8, §3.1]: A homomorphism  $X \xrightarrow{f} Y$  of  $p$ -compact groups is a pointed map  $BX \xrightarrow{Bf} BY$ . The homomorphism  $f$  is an isomorphism if  $Bf$  is a homotopy equivalence. It is a monomorphism if the homotopy fiber  $Y/X$  of  $Bf$  is  $\mathbb{F}_p$ -finite or equivalently if  $H^*(BX, \mathbb{F}_p)$  is a finitely generated module over  $H^*(BY, \mathbb{F}_p)$  via  $Bf^*$ .
- **Centralizers** [8, §3.4]: For a homomorphism  $Y \xrightarrow{f} X$  of  $p$ -compact groups, the centralizer  $C_X(f(Y))$  is defined by the equation  $BC_X(f(Y)) := \text{Map}(BY, BX)_{Bf}$ .
- **Maximal tori** [8, Definition 8.9]: A monomorphism  $T \hookrightarrow X$  of a  $p$ -compact torus into a  $p$ -compact group  $X$  is a maximal torus if  $C_X(T)$  is a  $p$ -compact toral group and if  $C_X(T)/T$  is homotopically discrete. Every  $p$  compact group admits maximal tori [8, Theorem 8.13].
- **Weyl group** [8, Definition 9.2]: Let  $BT_X \xrightarrow{Bf_T} BX$  be a maximal torus of a  $p$ -compact group  $X$ . Assume that  $Bf_T$  is already a fibration and treat  $\mathcal{W}_X$  as the space of self-maps of  $BT_X$  over  $BX$ . Composition gives  $\mathcal{W}_X$  the structure of an associative topological monoid. It is shown [8, Proposition 9.5] that  $\mathcal{W}_X$  is homotopically discrete and therefore  $W_X := \pi_0 \mathcal{W}_X$  is a (finite) group. Moreover, if  $X$  is connected, the action of  $W_X$  on  $BT_X$  induces a faithful representation

$$W_X \hookrightarrow \text{GL}(H_{\mathbb{Q}_p}^* BT_X) \cong \text{GL}_n(\mathbb{Q}_p^\wedge)$$

whose image is generated by pseudoreflections, i.e.  $W_X$  is a pseudo reflection group [8, Theorem 9.7].

- **Maximal torus normalizers** [8, Definition 9.8]: Let  $BT_X \xrightarrow{Bf_T} BX$  be a maximal torus of a  $p$ -compact group  $X$ . The normalizer of  $T_X$  denoted by  $NT_X$ , or simply by  $N_X$  or  $N$ , is the loop space such that  $BNT_X$  is the Borel construction associated to the action of  $\mathcal{W}_X$  on  $BT_X$ .

All these concepts generalize the classical algebraic definitions. In particular, if  $G$  is a compact Lie group such that  $\pi_0 G$  is  $p$ -group,  $i: T \longrightarrow G$  a maximal torus of  $G$ ,  $W$  the Weyl group of  $G$ , and  $N$  is the normalizer of the maximal torus  $T$ , then the  $p$ -completion  $i_p^\wedge: T_p^\wedge \longrightarrow G_p^\wedge$  is a maximal torus of the  $p$ -compact group  $G_p^\wedge$ . The Weyl group  $W$  is naturally isomorphic to the Weyl group  $W_{G_p^\wedge}$ . The classifying space  $BN$  of the normalizer  $N$  sits in the fibration  $BT \longrightarrow BN \longrightarrow BW$ , and a normalizer of the maximal torus  $T_p^\wedge$  of  $G_p^\wedge$  is isomorphic to the fiberwise  $p$ -completion  $BN_p^\circ$  by [23, Proposition 1.8], or [34, Lemma 6.1].

### 3. CONNECTEDNESS

In this section we proceed with the first step in the proof of Theorem 1.2 by proving the following proposition.

**Proposition 3.1.** *Let  $X$  be a 2-compact group with the normalizer of a maximal torus isomorphic to that of  $Sp(n)_2^\wedge$ , where  $n \geq 3$ , then  $X$  is connected.*

The proof of the result above requires calculating the Weyl group of some centralizer in the connected component of  $X$ . This is done by means of the technics developed by Dwyer and Wilkerson in [8] that we recall now.

An extended  $p$ -discrete torus  $P$  is an extension of a  $p$ -discrete torus  $(\mathbb{Z}/p^\infty)^n$  by a finite group. A discrete approximation for an extended  $p$ -compact torus  $P$  is a homomorphism  $f: \check{P} \longrightarrow P$ , where  $\check{P}$  is an extended  $p$ -discrete torus and  $f$  induces an isomorphism  $\check{P}/\check{P}_0 \longrightarrow \pi_0 P$  and an isomorphism  $H^* B\check{P}_0 \longrightarrow H^* B\check{P}_0$ . Every extended  $p$ -compact torus has a discrete approximation [8, Proposition 3.13].

**Definition 3.2.** [8, Definition 7.3] If  $s \in W$  is a pseudoreflection of order  $\text{ord}(s)$ , then

- (1) the *fixed point set*  $F(s)$  of  $s$  is the fixed point set of the action of  $x$  on  $\check{T}$  by conjugation, where  $x \in \check{N}(T)$  is an element which projects on  $s$  by natural projection  $\check{N}(T) \longrightarrow W$ ,
- (2) the *singular hyperplane*  $H(s)$  of  $s$  is the maximal divisible subgroup of  $F(s)$  (so  $H(s) \cong (\mathbb{Z}/p^\infty)^{r-1}$ ),
- (3) the *singular coset*  $K(s)$  of  $s$  is the subset of  $\check{T}$  given by elements of the form  $x^{\text{ord}(s)}$ , as  $x$  runs through elements of  $\check{N}(T)$ , which project to  $s$  in  $W$ , and
- (4) the *singular set*  $\sigma(s)$  of  $s$  is the union  $\sigma(s) = H(s) \cup K(s)$ .

Notice that there are inclusions  $H(s) \subset \sigma(s) \subset F(s)$  [8, Remark 7.7].

Let  $A \subset \check{T}$  be a subgroup. Let  $W_X(A)$  denote the Weyl group of  $C_X(A)$ , and  $W_X(A)_1$  the Weyl group of the unit component  $C_X(A)_0$  of  $C_X(A)$ . There are inclusions  $W_X(A)_1 \subset W_X(A) \subset W$ , where the last follows from [8, §4]. The next theorem tells how to calculate  $W_X(A)$  and  $W_X(A)_1$ .

**Theorem 3.3.** [8, Theorem 7.6] *Let  $X$  be a connected  $p$ -compact group with maximal torus  $T$  and Weyl group  $W$ . Suppose that  $A \subset \check{T}$  is a subgroup. Then*

- (1)  $W_X(A)$  is the subgroup of  $W$  consisting of the elements which, under the conjugation action of  $W$  on  $\check{T}$ , pointwise fix the subgroup  $A$ , and
- (2)  $W_X(A)_1$  is the subgroup of  $W_X(A)$  generated by those elements  $s \in W_X(A)$  such that  $s \in W$  is a reflection and  $A \subset \sigma(s)$ .

Now we have all the ingredients needed for the proof of Proposition 3.1

*Proof of Proposition 3.1.* Let  $X_0$  be the unit component of  $X$ , and let  $W_{X_0}$  be the Weyl group of  $X_0$ . Then  $W_{X_0}$  is a normal subgroup of  $W_X$  of index a power of 2 by [22, Proposition 3.8]. The minimal normal subgroup of  $W_X$  of 2 power index, usually denoted by  $O^2(W_X)$  in the literature, equals  $(\mathbb{Z}/2)^{n-1} \rtimes A_n$ , i.e. the sequence

$$(\mathbb{Z}/2)^{n-1} \rtimes A_n = O^2(W_X) \longrightarrow (\mathbb{Z}/2)^n \rtimes \Sigma_n \xrightarrow{\pi} (\mathbb{Z}/2)^2,$$

where  $A_n$  is the alternating group, is exact. The group  $(\mathbb{Z}/2)^2$  has 5 subgroups: the trivial subgroup 1, the first and the second factor  $Z_1$  and  $Z_2$ , the diagonal  $D$ , and the whole group  $(\mathbb{Z}/2)^2$ . Hence, there are 5 normal subgroups of  $W_X$  of index a power of 2:

- (1)  $\pi^{-1}(1) = (\mathbb{Z}/2)^{n-1} \rtimes A_n$ ,
- (2)  $\pi^{-1}(Z_1) = (\mathbb{Z}/2)^n \rtimes A_n$ ,
- (3)  $\pi^{-1}(Z_2) = (\mathbb{Z}/2)^{n-1} \rtimes \Sigma_n$ ,
- (4)  $\pi^{-1}(D)$ , and
- (5)  $\pi^{-1}((\mathbb{Z}/2)^2) = W_X$ .

Because  $W_{X_0}$  is the Weyl group of a connected 2-compact group,  $W_{X_0}$  is a pseudoreflection group. According to the Clark-Ewing list [5], only the cases (3) and (5) are pseudoreflection groups (note that  $n \geq 3$ ). We complete the proof showing that the case  $W_{X_0} = (\mathbb{Z}/2)^{n-1} \rtimes \Sigma_n$  is not possible.

Suppose that  $X$  is non connected, and let  $X_0$  be the unit component. By the arguments above  $W_{X_0}$  is  $(\mathbb{Z}/2)^{n-1} \rtimes \Sigma_n$ . Let  $V$  be the subgroup of the maximal torus  $T$  of  $X$  (and also of  $X_0$ ) generated by elements  $(-1, -1, 1, \dots)$ ,  $(1, 1, -1, -1, 1, \dots)$ , and so on. Then  $V$  is an elementary abelian 2-group of rank  $m = \lfloor \frac{n}{2} \rfloor$ . Let us write  $n = 2m + r$  where  $r$  is 0 or 1, and let  $C$  denote the centralizer  $C_{X_0}(V)$ . By Theorem 3.3(1), we get

$$W_C = \{s \in W_0 \mid s|_V = id_V\} = (\mathbb{Z}/2)^{n-1} \rtimes (\mathbb{Z}/2)^m,$$

where the subgroup  $(\mathbb{Z}/2)^m \subset \Sigma_n$  is generated by the transpositions  $\tau_{2i-1, 2i}$  for  $i = 1, \dots, m$ . Let  $C_0$  be the unit component of  $C$ . By Theorem 3.3(2), the Weyl group of  $C_0$  is

$$W_{C_0} = \langle s \in W_C \mid s \text{ is a reflection and } V \subset \sigma(s) \rangle.$$

An element  $s \in W_C = (\mathbb{Z}/2)^{n-1} \rtimes (\mathbb{Z}/2)^m$  is a reflection if and only if  $s$  equals  $((1, \dots, 1), \tau_{2i-1, 2i})$  or  $((1, \dots, 1, -1, -1, 1, \dots, 1), \tau_{2i-1, 2i})$  for some  $i$ , where both entries “-1” are in the  $(2i-1)^{\text{th}}$  and  $(2i)^{\text{th}}$  positions. We analyze both cases:

- If  $s = ((1, \dots, 1), \tau_{2i-1, 2i})$ , then  $F(s) = \{(x_1, \dots, x_n) \in (\mathbb{Z}/2^\infty)^n \mid x_{2i-1} = x_{2i}\}$  and  $H(s) = F(s)$ . Therefore  $\sigma(s) = F(s)$ .

- If  $s = ((1, \dots, 1, -1, -1, 1, \dots, 1), \tau_{2i-1, 2i})$ , then  $F(s) = \{(x_1, \dots, x_n) \in (\mathbb{Z}/2^\infty)^n \mid x_{2i-1} = x_{2i}^{-1}, i = 1, \dots, m\}$ . Hence also in this case  $H(s) = F(s) = \sigma(s)$ .

Since  $(-1)^{-1} = -1 \in \mathbb{Z}/2^\infty$ , the group  $V$  is a subgroup of  $\sigma(s)$  in both cases, and by Theorem 3.3, we get

$$W_{C_0} = \langle s \in W_C \mid s \text{ is a reflection} \rangle = ((\mathbb{Z}/2)^2)^m = (\mathbb{Z}/2)^{2m}.$$

Hence the normalizer of a maximal torus of  $C_0$  has the form  $M^m$ , where  $M$  is the subgroup of the normalizer of the maximal torus of  $Sp(2)_2^\wedge$  corresponding to the subgroup  $\langle ((1, 1), \tau_{1,2}), ((-1, -1), \tau_{1,2}) \rangle < (\mathbb{Z}/2)^2 \rtimes \mathbb{Z}/2 = W_{Sp(2)_2^\wedge}$ . By [12, Theorem 6.1] and [10, Theorem 0.5B (5)], the group 2-compact group  $C_0$  splits into a product  $C_0 \cong X_1 \times \dots \times X_m$ , where each  $X_i$  is isomorphic to  $(SU(2)^2/E_i)_2^\wedge$  for some subgroup  $E_i < Z(SU(2)^2) = (\mathbb{Z}/2)^2$ , and  $M$  is isomorphic to the maximal torus normalizer of  $X_i$ . Among the 5 possibilities for each  $X_i$ :

- (1)  $(SU(2) \times SU(2))_2^\wedge = Spin(4)_2^\wedge$ ,
- (2)  $(SU(2)/(\mathbb{Z}/2) \times SU(2))_2^\wedge \cong (SO(3) \times SU(2))_2^\wedge$ ,
- (3)  $(SU(2) \times SU(2)/(\mathbb{Z}/2))_2^\wedge \cong (SU(2) \times SO(3))_2^\wedge$ ,
- (4)  $(SU(2) \times_{\mathbb{Z}/2} SU(2))_2^\wedge \cong SO(4)_2^\wedge$ , and
- (5)  $((SU(2) \times SU(2))/(\mathbb{Z}/2)^2)_2^\wedge \cong (SO(3) \times SO(3))_2^\wedge$ ,

only  $SO(4)$  produces a pseudoreflection group which is equivalent to that given by  $M$ . But while the maximal torus normalizer of  $SO(4)$  is a split extension  $T : (\mathbb{Z}/2 \times \mathbb{Z}/2)$ ,  $M$  is not. Therefore there is no 2-compact group  $X_i$  whose maximal torus normalizer is  $M$ , what contradicts our initial assumption of  $X$  being non connected.  $\square$

#### 4. MOD-2 COHOMOLOGY OF THE 2-COMPACT GROUP $X$

In this section we calculate the mod-2 cohomology of a 2-compact group  $X$  whose maximal torus normalizer is isomorphic to that of  $Sp(n)_2^\wedge$ . This is done under the induction hypothesis that  $Sp(m)_2^\wedge$  is  $N$ -determined for  $2 < m < n$ . Notice that we already know that  $Sp(1)$  and  $Sp(2)$  are weakly  $N$ -determined.

First we need some information about the centralizers of elementary abelian subgroups in  $Sp(n)$ . It is well known that these centralizers are isomorphic to products  $Sp(n_1) \times \dots \times Sp(n_k)$ , where  $n_1 + \dots + n_k = n$ . The next lemma shows that those centralizers are  $N$ -determined if each factor is so.

**Lemma 4.1.** *Let  $Sp(m)_2^\wedge$  be an  $N$ -determined 2-compact group for  $2 < m \leq n$ . Then the product  $Sp(n_1)_2^\wedge \times \dots \times Sp(n_k)_2^\wedge$  is weakly  $N$ -determined if all  $n_i \leq n$  and is  $N$ -determined if  $2 < n_i \leq n$ .*

*Proof.* Let  $Y$  be a 2-compact group with maximal torus normalizer  $N_Y$  isomorphic to that of  $Sp(n_1)_2^\wedge \times \dots \times Sp(n_k)_2^\wedge$ . If at least one  $n_i$  is 1 or 2, assume that  $Y$  is connected. Since  $N_Y$  is a product  $N_1 \times \dots \times N_k$ , where  $N_i$  is the normalizer of a maximal torus of  $Sp(n_i)_2^\wedge$ , the space  $Y$  is by [12, Theorem 6.1] isomorphic to a product  $Y_1 \times \dots \times Y_k$ , where  $N_i$  is the normalizer of a maximal torus of  $Y_i$ . Hence  $Y_i$  is isomorphic to  $Sp(n_i)_2^\wedge$  and therefore  $Y$  is isomorphic to  $Sp(n_1)_2^\wedge \times \dots \times Sp(n_k)_2^\wedge$ . So  $Sp(n_1)_2^\wedge \times \dots \times Sp(n_k)_2^\wedge$  is weakly  $N$ -determined if at least one  $n_i$  is 1 or 2, otherwise  $Y$  is  $N$ -determined  $\square$

As  $X$  and  $Sp(n)_2^\wedge$  “share” the same maximal torus normalizer  $N$ , they both “share” the same maximal torus  $T$ . Let  $E_T < T$  be the maximal toral elementary abelian 2-group in both  $X$  and  $Sp(n)_2^\wedge$ . Call  $f_{E_T}$  the monomorphism  $E_T \hookrightarrow X$ . Next lemma shows that  $E_T$  is in fact the maximal elementary abelian subgroup of  $X$  (up to conjugation).

**Lemma 4.2.** *Let  $g: E \hookrightarrow X$  be an elementary abelian subgroup of  $X$ . Then  $g$  factors through  $f_{E_T}$ .*

*Proof.* If  $g: E \hookrightarrow X$  is central, then by [22, Lemma 4.1] or [8, Theorem 1.2]  $g$  factors through  $f_{E_T}$  (recall that  $X$  is connected by Proposition 3.1).

Now assume that  $g: E \hookrightarrow X$  is not central, thus there exists a subgroup  $V < E$  of rank 1 which is noncentral. By [18, Proof of Theorem 1.3] there exists  $\tilde{g}: E \hookrightarrow N$  such that  $Bg \simeq f_N B\tilde{g}$ , the centralizer  $C_N(\tilde{g})$  is the maximal torus normalizer of  $C_X(g)$ , and  $\tilde{g}|_V$  factors through  $f_{E_T}$ . Because  $V$  is a toral subgroup, the centralizer  $C_N(V)$  is the maximal torus normalizer of both  $C_{Sp(n)_2^\wedge}(V)$  and  $C_X(V)$  by [18, Theorem 1.3]. So the calculation of  $W_X(V)$  and  $W_X(V)_1$  by means of Theorem 3.3 equals the calculation of  $W_{Sp(n)_2^\wedge}(V)$  and  $W_{Sp(n)_2^\wedge}(V)_1$  what implies that  $C_X(V)$  is connected and since by induction, the centralizer  $C_{Sp(n)_2^\wedge}(V) = BSp(m)_2^\wedge \times BSp(n-m)_2^\wedge$ ,  $m > 0$ , is weakly  $N$ -determined (Lemma 4.1), then  $C_X(V)$  is isomorphic to  $C_{Sp(n)_2^\wedge}(V)$ .

The map  $g: E \hookrightarrow X$  has a lift to a map  $g': E \hookrightarrow C_V(X) \cong Sp(m)_2^\wedge \times Sp(n-m)_2^\wedge$ . Up to conjugacy every elementary abelian subgroup of  $Sp(m) \times Sp(n-m)$  is toral. Hence  $g$  is toral, i.e. factors through  $f_{E_T}$ .  $\square$

We can calculate the centralizer of  $E_T$  in  $X$ :

**Lemma 4.3.** *The centralizer  $C_X(E_T)$  is isomorphic to the 2-compact group  $(Sp(1)^n)_2^\wedge$ .*

*Proof.* As  $E_T$  is toral, the centralizer  $C_N(E_T)$  is the maximal torus normalizer of both  $C_{Sp(n)_2^\wedge}(E_T)$  and  $C_X(E_T)$  [19, Proposition 3.4(3)]. So the calculation of  $W_X(E_T)$  and  $W_X(E_T)_1$  by means of Theorem 3.3 equals the calculation of  $W_{Sp(n)_2^\wedge}(E_T)$  and  $W_{Sp(n)_2^\wedge}(E_T)_1$  what implies that  $C_X(E_T)$  is connected. Since  $C_{Sp(n)_2^\wedge}(E_T) = (C_{Sp(n)_2^\wedge}(E_T))_2^\wedge = (Sp(1)^n)_2^\wedge$  is weakly  $N$ -determined, the centralizer  $C_X(E_T)$  is isomorphic to  $C_{Sp(n)_2^\wedge}(E_T)$  by Lemma 4.1. Finally  $C_{Sp(n)_2^\wedge}(E_T) \cong (Sp(1)^n)_2^\wedge$ .  $\square$

The action of  $\Sigma_n < W_{Sp(n)} = W_X$  on  $BE_T$  induces an action of  $\Sigma_n$  on  $BC_X(E_T) = \text{Map}(BE_T, BX)_{Bf_{E_T}} \cong (Sp(1)_2^\wedge)^n$  that permutes the copies  $Sp(1)_2^\wedge$ . Define  $BY = BC_X(E_T) \times_{\Sigma_n} E\Sigma_n$  and consider the diagram

$$\begin{array}{ccccc}
 (BSp(1)^n)_2^\wedge & \xrightarrow{\quad} & BY & \xrightarrow{\quad} & B\Sigma_n \\
 \uparrow & & \uparrow & & \parallel \\
 (BT)_2^\wedge & \xrightarrow{\quad} & \text{Map}(BT, BX)_{Bf_T} \times_{\Sigma_n} E\Sigma_n & \xrightarrow{\quad} & B\Sigma_n \\
 \parallel & & \downarrow & & \downarrow \\
 (BT)_2^\wedge & \xrightarrow{\quad} & \text{Map}(BT, BX)_{Bf_T} \times_{W_{Sp(n)}} EW_{Sp(n)} & \xrightarrow{\quad} & BW_{Sp(n)}
 \end{array}$$

where all rows are fibrations. The space  $\text{Map}(BT, BX)_{Bf_T} \times_{W_{Sp(n)}} EW_{Sp(n)}$  is the normalizer of the maximal torus  $T$ , so it is isomorphic to  $B(T \rtimes W_{Sp(n)})$ . Therefore the space  $\text{Map}(BT, BX)_{Bf_T} \times_{\Sigma_n} E\Sigma_n$  is isomorphic to  $B(T \rtimes \Sigma_n)$ . Which means that the middle row has a section, and hence also the top row has a section. It follows that  $BY$  is homotopic to  $B((Sp(1)_2^n) \rtimes \Sigma_n)$ .

**Proposition 4.4.** *The cohomology  $H^*BX$  is detected by elementary abelian 2-subgroups.*

*Proof.* The cohomology  $H^*BSp(1)^n$  is detected by elementary abelian 2-subgroups, hence by [15],  $H^*BY$  is detected by elementary abelian subgroups. The normalizer  $Bf_N$  factors through the map  $Bf_Y$ . Because  $Bf_N^*$  is a monomorphism,  $Bf_Y^*$  is a monomorphism. Hence  $H^*BX$  is detected by elementary abelian 2-subgroups.  $\square$

We can now identify the algebra  $H^*BX$ :

**Proposition 4.5.** *The cohomology  $H^*BX$  is isomorphic to  $H^*BSp(n)$  as an algebra over the mod-2 Steenrod algebra.*

*Proof.* By Proposition 4.4, the cohomology  $H^*BX$  is detected by elementary abelian 2-subgroups, and by Lemma 4.2, every elementary abelian subgroup of  $X$  factors through  $E_T$ . Therefore  $H^*BX$  injects into  $H^*BE_T$  and therefore into  $H^*BC_X(E_T)$ . If we take trivial action of  $\Sigma_n$  on  $X$ , the inclusion  $C_X(E_T) \longrightarrow X$  is a  $\Sigma_n$ -equivariant map. Hence the cohomology  $H^*BX$  is a subalgebra of  $(H^*BSp(1)^n)^{\Sigma_n} = H^*BSp(n)$ . But  $H^*(BX; \mathbb{Q}) = H^*(BT; \mathbb{Q})^{W_X} = \mathbb{Q}[x_4, \dots, x_{4n}]$ , hence the Bockstein spectral sequence associated to  $H^*BX \subset H^*Sp(n) = \mathbb{F}_2[x_4, \dots, x_{4n}]$  converges to  $\mathbb{F}_2[x_4, \dots, x_{4n}]$ , and therefore  $H^*BX \cong H^*BSp(n)$ .  $\square$

Recall that the Quillen category  $Q_p(G)$  of a group  $G$  at a prime  $p$  is the category with objects  $(V, \alpha)$ , where  $V$  is a nontrivial elementary abelian  $p$ -group and  $\alpha: V \longrightarrow G$  is a monomorphism, and  $\text{Mor}_{Q_p(G)}((V, \alpha), (V', \alpha'))$  is the set of group morphisms  $f: V \longrightarrow V'$  such that  $\alpha = \alpha' \circ f$ . By Lannes' theory ([17]) and Dwyer-Zabrodsky theorem ([14], [29]), the set of monomorphisms  $\alpha: V \longrightarrow G$  is in bijection with the set of morphisms  $B\alpha^*: H^*BG \longrightarrow H^*BV$  of unstable algebras over the Steenrod algebra  $\mathcal{A}_p$  such that  $H^*BV$  is a finitely generated module over  $B\alpha^*(H^*BG)$ . Hence, there is an equivalent description of the Quillen category which can be used also for  $p$ -compact groups ([13, §2]). If  $X$  is a  $p$ -compact group, then  $Q_p(X)$  is the category with objects  $(V, \alpha)$ , where  $V$  is a nontrivial elementary abelian  $p$ -group and  $\alpha: H^*BX \longrightarrow H^*BV$  is a monomorphism of unstable algebras over the Steenrod algebra  $\mathcal{A}_p$  such that  $H^*BV$  is a finitely generated module over  $\alpha^*(H^*BX)$ , and  $\text{Mor}_{Q_p(G)}((V, \alpha), (V', \alpha'))$  is a set of group morphisms  $f: V \longrightarrow V'$  such that  $\alpha = Bf^*\alpha'$ . If  $X$  is a compact Lie group then the definitions agree ([13, Proposition 2.2]).

By the definition of the Quillen category and propositions 4.4 we get the following proposition.

**Proposition 4.6.** *The categories  $Q_2(Sp(n))$  and  $Q_2(X)$  are isomorphic.*



5. 2-STUBBORN DECOMPOSITION OF  $Sp(n)$ 

A 2-stubborn subgroup of a Lie group  $G$  is a 2-toral group  $P$  such that  $N_G(P)/P$  is a finite group which has no nontrivial normal 2-subgroup. Let  $\mathcal{R}_2(Sp(n))$  be the 2-stubborn category of  $Sp(n)$ , which is the full subcategory of the orbit category of  $Sp(n)$  with objects  $Sp(n)/P$ , where  $P \subset Sp(n)$  is a 2-stubborn subgroup. The natural map

$$\operatorname{hocolim}_{Sp(n)/P \in \mathcal{R}_2(Sp(n))} ESp(n)/P \longrightarrow BSp(n)$$

induces an isomorphism of homology with  $\mathbb{Z}_{(2)}$ -coefficients [16, Theorem 4]. Hence, defining a map  $f: Sp(n)_2^\wedge \longrightarrow X$  is equivalent to defining a family of compatible maps  $\{f_P: ESp(n)/P \simeq BP \longrightarrow X \mid Sp(n)/P \in \operatorname{ob}(\mathcal{R}_2(Sp(n)))\}$ .

We first recall the 2-stubborn subgroups of  $Sp(n)$  calculated in [31]. Let the permutations  $\sigma_0, \dots, \sigma_{k-1}$  in  $\Sigma_{2^k}$  be defined by

$$\sigma_r(s) = \begin{cases} s + 2^r; & s \equiv 1, \dots, 2^r \pmod{2^{r+1}} \\ s - 2^r; & s \equiv 2^r + 1, \dots, 2^{r+1} \pmod{2^{r+1}} \end{cases}$$

Let  $A_0, \dots, A_{k-1} \in Sp(2^k)$  be diagonal matrices with

$$(A_r)_{ss} = (-1)^{\lfloor \frac{s-1}{2^r} \rfloor},$$

where  $\lfloor - \rfloor$  denotes greatest interger, and  $B_0, \dots, B_{k-1}$  be the permutation matrices for the  $\sigma_0, \dots, \sigma_{k-1}$ .

**Definition 5.1.** For every  $k \geq 0$ , the subgroups  $E_{2^k} \subset \Sigma_{2^k}$  and  $\Gamma_{2^k}, \bar{\Gamma}_{2^k} \subset Sp(2^k)$  are defined by

$$\begin{aligned} E_{2^k} &= \langle \sigma_0, \dots, \sigma_{k-1} \rangle \cong (\mathbb{Z}/2)^k, \\ \Gamma_{2^k} &= \langle uI, A_r, B_r \mid u \in Q(8), 0 \leq r < k \rangle, \\ \bar{\Gamma}_{2^k} &= \langle uI, A_r, B_r \mid u \in S^1(j), 0 \leq r < k \rangle, \end{aligned}$$

where  $Q(8) = \{\pm 1, \pm i, \pm j, \pm k\}$  is the quaternion group and  $S^1(j) = \{a + bi, aj + bk \mid a^2 + b^2 = 1\}$  is the normalizer of the maximal torus in  $Sp(1) = S^3$ .

**Remark 5.2.** Let  $P$  be  $\Gamma_{2^k}$  or  $\bar{\Gamma}_{2^k}$  and  $P_D$  subgroup of all diagonal matrices in  $P$ . Then  $P_D$  is  $Q(8) \times E_{2^k}$  or  $S^1(j) \times E_{2^k}$  and the exstension  $P_D \longrightarrow P \longrightarrow (\mathbb{Z}/2)^k$  splits.

**Theorem 5.3.** [31, Theorem 3]

(1) A 2-stubborn group  $P < Sp(n)$  is irreducible if it is conjugate to either

$$P = \Gamma_{2^k} \wr E_{2^{r_1}} \wr \dots \wr E_{2^{r_s}} \text{ or } P = \bar{\Gamma}_{2^k} \wr E_{2^{r_1}} \wr \dots \wr E_{2^{r_s}}$$

where  $n = 2^{k+r_1+\dots+r_s}$ .

(2) A group  $P < Sp(n)$  is a 2-stubborn group if it is conjugate to  $P_1 \times \dots \times P_s$ , where  $P_i$  is an irreducible 2-stubborn subgroup of  $Sp(n_i)$  and  $n = n_1 + \dots + n_s$ .

Let  $\tilde{\mathcal{R}}_2(Sp(n))$  be the full subcategory of  $\mathcal{R}_2(Sp(n))$  with objects  $Sp(n)/P$ , where  $P$  is one of the representative 2-stubborn groups from the previous theorem. The category  $\tilde{\mathcal{R}}_2(Sp(n))$  is equivalent to  $\mathcal{R}_2(Sp(n))$ , so the natural map

$$\operatorname{hocolim}_{Sp(n)/P \in \tilde{\mathcal{R}}_2(Sp(n))} ESp(n)/P \longrightarrow BSp(n)$$

is also a homotopy equivalence up to 2-completion.

**Proposition 5.4.** *Let  $Sp(n)/P \in \tilde{\mathcal{R}}_2(Sp(n))$  and define  $P_D = P \cap Sp(1)^n$  and  $P_T = P \cap T_{Sp(n)}$ . Then*

- (1)  $C_{Sp(n)}(P_T) = T_{Sp(n)}$  and  $C_{Sp(n)}(P_D) = (\mathbb{Z}/2)^n$ ,
- (2) *for any extension  $\alpha: P \longrightarrow Sp(n)$  of  $i: P_T \longrightarrow Sp(n)$ , we have  $C_{Sp(n)}(\alpha(P)) = Z(P)$  and*
- (3) *the canonical map*

$$\pi_0(\operatorname{Map}(BP, BSp(n)_2^\wedge)_{B\alpha|_{BP_T} = Bi_{P_T}}) \longrightarrow \operatorname{Hom}(H^*BSp(n), H^*BP)$$

*is an injection.*

**Remark 5.5.** By  $\operatorname{Map}(BP, BSp(n)_2^\wedge)_{B\alpha|_{BP_T} = Bi_{P_T}}$  we denote the components of the mapping space  $\operatorname{Map}(BP, BSp(n)_2^\wedge)$  given by maps  $B\alpha: BP \longrightarrow Sp(n)_2^\wedge$ , such that  $B\alpha|_{BP_T} \simeq Bi_{P_T}$ .

*Proof.* Part (1) is obvious for  $P = \Gamma_{2^k}$  and  $P = \bar{\Gamma}_{2^k}$ . If  $P = Q \wr E_{2^r}$ , where  $Q$  is irreducible 2-stubborn subgroup of  $Sp(2^{k-r})$ , then  $C_{Sp(2^k)}(P_T) = C_{Sp(2^{k-r})}(Q_T)^{2^r}$  which is, by induction,  $(T_{Sp(2^{k-r})})^{2^r} = T_{Sp(2^k)}$ . If  $P = P_1 \times \dots \times P_s$  is a product of irreducible 2-stubborn groups, then  $C_{Sp(n)}(P_T) = C_{Sp(n_1)}((P_1)_T) \times \dots \times C_{Sp(n_s)}((P_s)_T) = T_{Sp(n_1)} \times \dots \times T_{Sp(n_s)} = T_{Sp(n)}$ . Analogously we prove that  $C_{Sp(n)}(P_D) = (\mathbb{Z}/2)^n$ .

Every map  $g \in \operatorname{Map}(BP, BSp(n)_2^\wedge)$  is homotopic to  $B\alpha: BP \longrightarrow BSp(n)_2^\wedge$ , where  $\alpha$  is the 2-completion of a homomorphism of groups  $P \longrightarrow Sp(n)$  ([14], [29]).

Let  $P$  be an irreducible 2-stubborn subgroup of  $Sp(2^k)$  and let  $\alpha: P \longrightarrow Sp(2^k)$  be a homomorphism such that  $\alpha|_{P_T} = i_{P_T}$ . The extensions  $B\alpha|_{BP_D}: BP_D \longrightarrow BSp(2^k)_2^\wedge$  of  $Bi_{P_T}$  are classified by obstruction groups  $H^m(P_D/P_T; \pi_m(\operatorname{Map}(BP_T, BSp(2^k)_2^\wedge)_{Bi_{P_T}}))$ . By [16],  $\operatorname{Map}(BP_T, BSp(2^k)_2^\wedge)_{Bi_{P_T}}$  is homotopy equivalent to  $BC_{Sp(2^k)}(P_T)_2^\wedge$ , which is isomorphic to  $(BS^1)^{2^k}$ , by part (1). Then

$$H^m(P_D/P_T; \pi_m(\operatorname{Map}(BP_T, BSp(2^k)_2^\wedge)_{Bi_{P_T}})) = H^m(P_D/P_T; \pi_m(BS^1)^{2^k})$$

and the only possible nontrivial group is for  $m = 1$ . And

$$H^1(P_D/P_T; \pi_m(\operatorname{Map}(BP_T, BSp(2^k)_2^\wedge)_{Bi_{P_T}})) = H^1(\mathbb{Z}/2; (\mathbb{Z}_2^\wedge)^{2^k}),$$

where the group  $\mathbb{Z}/2$  acts on  $(\mathbb{Z}_2^\wedge)^{2^k}$  by reflection on each component; this action comes from the action of the Weyl group  $\mathbb{Z}/2$  of the group  $Sp(1)$  on the maximal torus  $S^1$ . By Shapiro's lemma [3, III, Proposition 6.2], the group  $H^1(\mathbb{Z}/2; (\mathbb{Z}_2^\wedge)^{2^k})$  is trivial, so all obstruction groups vanish. Hence if  $B\alpha|_{BP_T} = Bi_{P_T}$  then  $B\alpha|_{BP_D} = Bi_{P_D}$ .

First we will prove part (2) and (3) for the case  $P$  is  $\Gamma_{2^k}$  or  $\overline{\Gamma}_{2^k}$ . Let  $\alpha: P \longrightarrow Sp(2^k)$  be a homomorphism such that  $\alpha|_{P_T} = i_{P_T}$ . Then by above paragraph  $B\alpha|_{BP_D}$  is homotopic to  $Bi_{P_D}$ . The extensions  $B\alpha: BP \longrightarrow BSp(2^k)_2^\wedge$  of  $Bi_{P_D}$  are classified by obstruction groups  $H^m(P/P_D; \pi_m(\text{Map}(BP_D, BSp(2^k)_2^\wedge)_{Bi_{P_D}}))$ . By [16], the mapping space  $\text{Map}(BP_D, BSp(2^k)_2^\wedge)_{Bi_{P_D}}$  is homotopy equivalent to  $BC_{Sp(2^k)}(P_D)_2^\wedge$ , which is isomorphic to  $(B\mathbb{Z}/2)^{2^k}$  (part (1)). Then the obstruction group

$$H^m(P/P_D; \pi_m(\text{Map}(BP_D, BSp(2^k)_2^\wedge)_{Bi_{P_D}})) = H^m(P/P_D; \pi_m(B(\mathbb{Z}/2)^{2^k}))$$

is nontrivial only possibly for  $m = 1$ . The group  $P/P_D$  is isomorphic to the group generated by the permutation matrices  $B_0, \dots, B_{k-1}$  (Definition 5.1). So  $P/P_D = (\mathbb{Z}/2)^k$  and the action of  $P/P_D$  on  $\pi_1(B(\mathbb{Z}/2)^{2^k}) = (\mathbb{Z}/2)^{2^k}$  is given by permutations  $\sigma_0, \dots, \sigma_{k-1}$  which define matrices  $B_0, \dots, B_{k-1}$ . By Shapiro's lemma [3, III, Proposition 6.2],

$$H^1(P/P_D; \pi_1(\text{Map}(BP_D, BSp(2^k)_2^\wedge)_{Bi_{P_D}})) = H^1(E_{2^k}; (\mathbb{Z}/2)^{2^k}) = H^1(1; \mathbb{Z}/2) = 1,$$

so all obstruction groups vanish. Therefore  $B\alpha$  is homotopic to  $Bi_P$  and  $C_{Sp(n)}(\alpha)$  equals  $Z(P)$ .

Now we will prove part (2) and (3) for an irreducible 2-stubborn subgroup  $P$  of  $Sp(2^k)$ . Let us write  $P = Q \wr E_{2^r}$ , where  $Q$  is an irreducible 2-stubborn subgroup of  $Sp(2^{k-r})$ . Let  $\alpha, \beta: P \longrightarrow Sp(2^k)$  be two homomorphisms such that  $B\alpha^* = B\beta^*$  and  $\alpha|_{BP_T} = i_{P_T} = \beta|_{BP_T}$ . We proved that  $\alpha|_{BP_D} = i_{P_D} = \beta|_{BP_D}$ . Let  $\bar{\alpha}, \bar{\beta}: Q^{2^r} \longrightarrow Sp(2^k)$  be the restrictions of  $\alpha$  and  $\beta$ . Because  $Z(Q^{2^r}) = Z(Q)^{2^r} = (\mathbb{Z}/2)^{2^r}$ , the homomorphisms  $\bar{\alpha}$  and  $\bar{\beta}$  factor through homomorphisms

$$\tilde{\alpha}, \tilde{\beta}: Q^{2^r} \longrightarrow C_{Sp(2^k)}(Z(Q^{2^r})) = Sp(2^{k-r})^{2^r}.$$

The map  $B\tilde{\alpha}$  is homotopic to the map

$$BQ^{2^r} \simeq \text{Map}(BZ(Q^{2^r}), BQ^{2^r})_{Bi} \xrightarrow{\text{Map}(BZ(Q^{2^r}), B\tilde{\alpha})} \text{Map}(BZ(Q^{2^r}), BSp(2^k)_2^\wedge)_{Bi} \simeq (BSp(2^{k-r})^{2^r})_2^\wedge,$$

hence  $H^*(B\tilde{\alpha}; \mathbb{F}_2) = H^*(\text{Map}(BZ(Q^{2^r}), B\tilde{\alpha}); \mathbb{F}_2)$ . Analogously  $H^*(B\tilde{\beta}; \mathbb{F}_2) = H^*(\text{Map}(BZ(Q^{2^r}), B\tilde{\beta}); \mathbb{F}_2)$ . By Lannes' theory [17],

$$H^*(\text{Map}(BZ(Q^{2^r}), B\tilde{\alpha}); \mathbb{F}_2) = T_{B\tilde{\alpha}^*}^{Z(Q^{2^r})} = T_{B\tilde{\beta}^*}^{Z(Q^{2^r})} = H^*(\text{Map}(BZ(Q^{2^r}), B\tilde{\beta}); \mathbb{F}_2),$$

so  $B\tilde{\alpha}^* = B\tilde{\beta}^*$ .

The homomorphisms  $\tilde{\alpha}$  and  $\tilde{\beta}$  are matrices of dimension  $2^r \times 2^r$ , where entries are  $\tilde{\alpha}_{i,j}, \tilde{\beta}_{i,j}: Q_i \longrightarrow Sp(2^{k-r})_j$ . The indexes  $i$  and  $j$  indicate the components in the products. By induction,  $B\tilde{\alpha}_{i,i}$  and  $B\tilde{\beta}_{i,i}$  are homotopic and therefore  $\tilde{\alpha}_{i,i}$  and  $\tilde{\beta}_{i,i}$  are conjugate [17, Théorème 3.4.5]. We can assume that  $\tilde{\alpha}_{i,i} = \tilde{\beta}_{i,i}$ . Because  $Q_i$  and  $Q_j$  commutes for  $i \neq j$ , the homomorphisms  $\tilde{\alpha}_{i,j}$  and  $\tilde{\beta}_{i,j}$  factor through homomorphisms  $\hat{\alpha}_{i,j}, \hat{\beta}_{i,j}: Q_i \longrightarrow C_{Sp(2^{k-r})}(\tilde{\alpha}_{j,j}(Q))$ . By induction, the centralizer  $C_{Sp(2^{k-r})}(\tilde{\alpha}_{j,j}(Q))$  equals  $Z(Q_j) = (\mathbb{Z}/2)_j$ . Because  $\tilde{\alpha}|_{P_D} = \tilde{\beta}|_{P_D}$ , the homomorphism

$\tilde{\alpha}_{i,j} \cdot \tilde{\beta}_{i,j}^{-1}: Q_i \longrightarrow (\mathbb{Z}/2)_j$  factors through a homomorphism  $\gamma_{i,j}: (Q/Q_D)_i \longrightarrow (\mathbb{Z}/2)_j$ . Then  $\tilde{\beta}_{i,j}$  equals to composition

$$Q_i \xrightarrow{\Delta} Q_i \times (Q/Q_D)_i \xrightarrow{\tilde{\alpha}_{i,j} \times \gamma_{i,j}} Sp(2^{k-r})_j \times (\mathbb{Z}/2)_j \xrightarrow{\mu} Sp(2^{k-r})_j,$$

where  $\Delta$  is the diagonal map composed by the quotient map and  $\mu$  is the multiplication in  $Sp(2^{k-r})$ . Because  $B\tilde{\alpha}_{i,j}^* = B\tilde{\beta}_{i,j}^*$ , the map  $B\gamma_{i,j}$  induces trivial map on mod-2 cohomology. Because  $Q/Q_D$  is an iterated wreath product of elementary abelian groups, the map  $\gamma_{i,j}$  is constant [24, Lemma 6.10]. Hence  $\tilde{\alpha}_{i,j} = \tilde{\beta}_{i,j}$  and so  $\bar{\alpha} = \bar{\beta}$  and the centralizer  $C_{Sp(2^k)}(\alpha)$  is given by the fixed-point set  $C_{Sp(2^k)}(\alpha) = (C_{Sp(2^{k-r})}(\tilde{\alpha}))^{E_{2^r}} = ((C_{Sp(2^{k-r})}(Q))^{2^r})^{E_{2^r}} = ((\mathbb{Z}/2)^{2^r})^{E_{2^r}} = \mathbb{Z}/2 = Z(P)$ , which proves part (2).

The extensions  $B\alpha: BP \longrightarrow BSp(2^k)_2^\wedge$  of  $B\bar{\alpha}$  are classified by the obstruction groups  $H^m(P/Q^{2^r}; \pi_m(\text{Map}(BQ^{2^r}, BSp(2^k)_2^\wedge)_{B\bar{\alpha}}))$ . By [16], the mapping space  $\text{Map}(BQ^{2^r}, BSp(2^k)_2^\wedge)_{B\bar{\alpha}}$  is homotopy equivalent to  $BC_{Sp(2^k)}(Q^{2^r})_2^\wedge = (\mathbb{Z}/2)$ . Hence the obstruction groups are

$$H^m(P/Q^{2^r}; \pi_m(\text{Map}(BQ^{2^r}, BSp(2^k)_2^\wedge)_{B\bar{\alpha}})) = H^m(P/Q^{2^r}; \pi_m(B\mathbb{Z}/2)).$$

The only possible nontrivial obstruction group is for  $m = 1$ . The group  $P/Q^{2^r} = E_{2^r}$  acts by permutation on  $BC_{Sp(2^n)}(Q^{2^r}) = (B\mathbb{Z}/2)^{2^r}$ , hence  $\text{Ind}_1^{E_{2^r}}(\mathbb{Z}/2)^{2^r} = (\mathbb{Z}/2)$ , by Shapiro's lemma [3, III, Proposition 6.2]. Therefore

$$H^1(K \wr E_{2^r}; (\mathbb{Z}/2)^{2^k}) = H^1(K; (\mathbb{Z}/2)^{2^{k-r}}).$$

Therefore all obstruction groups vanish, so  $B\alpha$  is homotopic to  $Bi_P$ .

Finally let  $P = P_1 \times \dots \times P_s$ , where  $P_i$  is an irreducible 2-stubborn subgroup of  $Sp(n_i)$ . Let  $\alpha, \beta: P \longrightarrow Sp(n)$  be two homomorphisms such that  $B\alpha^* = B\beta^*$  and  $\alpha|_{P_T} = \beta|_{P_T}$ . Both homomorphisms factor through  $\bar{\alpha}, \bar{\beta}: P \longrightarrow C_{Sp(n)}(Z(P)) = C_{Sp(n)}((\mathbb{Z}/2)^s) = Sp(n_1) \times \dots \times Sp(n_s)$ . In the same way as in the case  $P$  is irreducible 2-stubborn group, we can show that  $B\bar{\alpha}^* = B\bar{\beta}^*$ . Maps  $\bar{\alpha}$  and  $\bar{\beta}$  are matrices of dimension  $s \times s$  with entries maps  $\bar{\alpha}_{i,j}, \bar{\beta}_{i,j}: P_i \longrightarrow Sp(n_j)$ . Analogously as before we can show that  $\bar{\alpha}_{i,j} = \bar{\beta}_{i,j}$ , so  $B\alpha \simeq B\beta$ . The equation  $Z(P) = Z(P_1) \times \dots \times Z(P_s)$  finishes the proof.  $\square$

## 6. THE MAP FROM $Sp(n)_2^\wedge$ TO $X$

For every object  $Sp(n)/P$  in  $\tilde{\mathcal{R}}(Sp(n))$  we define a map  $f_P: P \longrightarrow X$  as the composition of the two inclusions  $i_P: P \longrightarrow N$  and  $f_N: N \longrightarrow X$ . We will prove that for every morphism  $c_g: Sp(n)/P \longrightarrow Sp(n)/Q$  in  $\tilde{\mathcal{R}}(Sp(n))$ , the diagram

$$(1) \quad \begin{array}{ccccc} BP & \xrightarrow{Bi_P} & BN & \xrightarrow{Bf_N} & BX \\ Bc_g \downarrow & & & & \parallel \\ BQ & \xrightarrow{Bi_Q} & BN & \xrightarrow{Bf_N} & BX \end{array}$$

commutes up to homotopy.

Let us define  $\alpha = f_N \circ i_P$  and  $\beta = f_N \circ i_Q \circ c_g$ . Then  $B\alpha^* = B\alpha^*$ . The group  $P_T = P \cap Sp(n)$  is 2-toral. The restrictions  $\alpha|_{P_T}$  and  $\beta|_{P_T}$  are conjugate in  $Sp(n)$ , and hence by [23, Proposition 4.1], they are also conjugate in the normalizer  $N$  of the maximal torus. So  $B\alpha|_{BP_T} \simeq B\beta|_{BP_T}$ . By the next proposition,  $B\alpha \simeq B\beta$ .

Let  $K \longrightarrow G \longrightarrow H$  be an exact sequence of groups. Then  $H$  acts freely on  $\widetilde{BK} = EG/K \simeq BK$ , and  $\widetilde{BK}/H$  equals  $BG$ . For any space  $BX$  with trivial action of the group  $H$ , we have

$$(2) \quad \begin{aligned} \text{Map}(BG, BX) &= \text{Map}(\widetilde{BK}/H, BX) = \text{Map}_H(\widetilde{BK}, BX) \simeq \\ &\simeq \text{Map}_H(EH \times \widetilde{BK}, BX) = \text{Map}_H(EH, \text{Map}(\widetilde{BK}, BX)) = \text{Map}(\widetilde{BK}, BX)^{hH}. \end{aligned}$$

**Proposition 6.1.** *For every  $Sp(n)/P \in \text{ob}(\widetilde{\mathcal{R}}(Sp(n)))$ , the canonical map*

$$\pi_0(\text{Map}(BP, BX)_{B\alpha|_{BP_T}=Bf_{P_T}}) \longrightarrow \text{Hom}(H^*BX, H^*BP)$$

*is an injection.*

*Proof.* Consider the diagram

$$\begin{array}{ccc} & \text{Map}(\widetilde{BP_T}, BY)_{Bi_T} & \\ \swarrow & & \searrow \\ \text{Map}(\widetilde{BP_T}, BSp(n)_2^\wedge)_{Bi_T} & & \text{Map}(\widetilde{BP_T}, BX)_{Bf_T}. \end{array}$$

By [29], the mapping space  $\text{Map}(\widetilde{BP_T}, BSp(n)_2^\wedge)_{Bi_T}$  is homotopy equivalent to  $BC_{Sp(n)}(BP_T)_2^\wedge$  and by proposition 5.4, this space is homotopy equivalent to  $(BT_{Sp(n)})_2^\wedge$ . Analogously  $\text{Map}(\widetilde{BP_T}, BY)_{Bi_T}$  is homotopy equivalent to  $(BT_{Sp(n)})_2^\wedge$ . The mapping space  $\text{Map}(\widetilde{BP_T}, BX)_{Bf_T}$  is the classifying space of a 2-compact group ([9]). Its Weyl group is  $\text{Iso}(Bf_N \circ Bi_{P_T}) = \{w \in W_X \mid w \circ Bf_N \circ Bi_{P_T} \simeq Bf_N \circ Bi_{P_T}\}$  ([34, Proposition 4.3]). By the construction of the map  $f_N$ , the group  $\text{Iso}(Bf_N \circ Bi_{P_T})$  equals  $\text{Iso}(Bi_N \circ Bi_{P_T})$ . Because  $\text{Iso}(Bi_N \circ Bi_{P_T})$  is the Weyl group of the mapping space  $\text{Map}(\widetilde{BP_T}, BSp(n)_2^\wedge)_{Bi_T} \simeq (BT_{Sp(n)})_2^\wedge$ , the group  $\text{Iso}(Bf_N \circ Bi_{P_T})$  is trivial, hence  $\text{Map}(\widetilde{BP_T}, BX)_{Bf_T} \simeq (BT_{Sp(n)})_2^\wedge$ . Therefore both maps in the diagram (6) are homotopy equivalences.

Taking homotopy fixed points we obtain the following diagram

$$\begin{array}{ccc} & \text{Map}(\widetilde{BP_T}, BY)_{Bi_T}^{h(P/P_T)} & \\ \swarrow & & \searrow \\ \text{Map}(\widetilde{BP_T}, BSp(n)_2^\wedge)_{Bi_T}^{h(P/P_T)} & & \text{Map}(\widetilde{BP_T}, BX)_{Bf_T}^{h(P/P_T)}, \end{array}$$

where both maps are mod-2 equivalences, since an equivariant mod-2 equivalence between 1-connected spaces induces a mod-2 equivalence between the homotopy fixed-point sets.

By proposition 5.4, the components of  $\text{Map}(\widetilde{BP_T}, BSp(n)_2^\wedge)_{Bi_T}^{h(P/P_T)}$  are distinguished by mod 2 cohomology. Any map in  $\text{Map}(\widetilde{BP_T}, BX)_{Bf_T}^{h(P/P_T)}$  has a lift to  $BN$  and therefore to  $BY$ . The obstruction group which classifies the extensions is

$$H^2(P/P_T; \pi_2 \text{Map}(\widetilde{BP_T}, BX)_{Bf_T}) \cong H^2(P/P_T; \pi_2 \text{Map}(\widetilde{BP_T}, BSp(n)_2^\wedge)_{Bi_T}),$$

so the components of  $\text{Map}(\widetilde{BP_T}, BX)_{Bf_T}^{h(P/P_T)} \simeq \text{Map}(BP, BX)_{B\alpha|_{BP_T}=Bi_{P_T}}$  are also distinguished by mod-2 cohomology.  $\square$

Diagram (1) establishes a map from the 1-skeleton of the homotopy colimit  $\{BP\}_{\widetilde{\mathcal{R}}_2(Sp(n))}$  to  $BX$ . The obstruction groups for extending a map defined on the 1-skeleton of the homotopy colimit to a map on the total homotopy colimit are

$$\varprojlim_{\widetilde{\mathcal{R}}_2(Sp(n))}^{i+1} \pi_i \text{Map}(BP, BX)_{Bf_P}$$

for  $i \geq 2$ , where  $\varprojlim^i$  is the  $i$ -th derived functor of the inverse limit functor ([4] and [36]).

Let  $\mathcal{A}b$  be the category of Abelian groups and let

$$\Pi_j^X, \Pi_j^{Sp(n)}: \widetilde{\mathcal{R}}_2(Sp(n)) \longrightarrow \mathcal{A}b$$

be functors defined by

$$\Pi_j^X(Sp(n)/P) = \pi_j \text{Map}(BP, BX)_{Bf_P},$$

$$\Pi_j^{Sp(n)}(Sp(n)/P) = \pi_j \text{Map}(BP, BSp(n)_2^\wedge)_{Bi_P}.$$

Note that  $\text{Map}(BP, BSp(n)_2^\wedge)_{Bi_P}$  is isomorphic to  $BZ(P)_2^\wedge$  ([16], Theorem 3.2) therefore  $\Pi_1(Sp(n))(Sp(n)/P)$  is well defined. By the next proposition, also  $\Pi_1(X)(Sp(n)/P)$  is well defined.

**Proposition 6.2.** *There exists a natural transformation*

$$\mathcal{T}: \Pi_j^{Sp(n)} \rightarrow \Pi_j^X$$

*which is an equivalence.*

*Proof.* For every 2-stubborn group  $P$  we have homotopy equivalences

$$(3) \quad \text{Map}(BP, BSp(n)_2^\wedge)_{Bi_P} \xleftarrow{\simeq} \text{Map}(BP, BY)_{Bi_P} \xrightarrow{\simeq} \text{Map}(BP, BX)_{Bf_P}$$

which depend on the chosen lift  $Bi_P: BP \longrightarrow BY$  of the map  $Bi_P: BP \longrightarrow BSp(n)_2^\wedge$ . Because  $\text{Rep}(P, Sp(n)) \longrightarrow [BP, BSp(n)]$  is a bijection ([14], [29]), two lifts differ by a conjugation  $Bc_g$ . Since  $Bf_P \simeq Bf_P \circ Bc_g$ , the equivalence (3) induces well defined isomorphisms

$$\Pi_j^{Sp(n)}(Sp(n)/P) \rightarrow \Pi_j^X(Sp(n)/P)$$

which commute with maps induced by morphisms in  $\widetilde{\mathcal{R}}_2(Sp(n))$ .  $\square$

**Proposition 6.3.** *For all  $i, j \geq 1$ ,*

$$\varprojlim_{\widetilde{\mathcal{R}}_2(Sp(n))}^i \pi_j \text{Map}(BP, BX)_{Bf_P} = 0.$$

*Proof.* By the previous lemma,

$$\varprojlim_{\mathcal{R}_2(Sp(n))}^i \pi_j \operatorname{Map}(BP, BX)_{Bf_P} = \varprojlim_{\mathcal{R}_2(Sp(n))}^i \pi_j \operatorname{Map}(BP, BSp(n)_2^\wedge)_{(Bi_P)_2^\wedge}$$

and the right side is 0 [16, Theorem 4.8].  $\square$

Because all obstructions vanish, there exists a map

$$f: \operatorname{hocolim}_{\mathcal{R}_2(Sp(n))} BP \longrightarrow BX.$$

By the construction of the map we have a commutative diagram

$$\begin{array}{ccc} & BY & \\ \swarrow & & \searrow \\ \operatorname{hocolim}_{\mathcal{R}_2(Sp(n))} ESp(n)/P & \xrightarrow{f} & BX \end{array}$$

where the diagonal maps induce monomorphisms on the cohomology and therefore  $f^*$  is a monomorphism. Since  $H^*BSp(n) \cong H^*BX$ ,  $f^*$  is an isomorphism and therefore  $f$  is a homotopy equivalence.

## 7. $Sp(n)$ AS A LOOP SPACE

The normalizer conjecture can be stated also for finite loop spaces with maximal torus normalizers as a weak version of Wilkerson' conjecture (see [35]).

**Theorem 7.1.** *Let  $L$  be a connected finite loop space with a maximal torus normalizer isomorphic to that of  $Sp(n)$ . Then  $BL$  is homotopy equivalent to  $BSp(n)$ .*

*Proof.* To prove  $BL \simeq BSp(n)$  is equivalent to showing that  $BL$  and  $BSp(n)$  lie in the same adic genus ([30]). The loop spaces  $BL$  and  $BSp(n)$  have the same rational genus. Since  $BL$  is finite and connected,  $L_p^\wedge$  is a  $p$ -compact group. The maximal torus normalizer of  $L_p^\wedge$  is just the fibrewise  $p$ -completion of  $N$  by the fibration  $BT \longrightarrow BN \longrightarrow BW_L$ . Hence  $L_p^\wedge$  and  $Sp(n)_p^\wedge$  have isomorphic normalizers of the maximal torus. By [2],  $BSp(n)_p^\wedge$  is  $N$ -determined if  $p$  is an odd prime and by the main theorem of this paper  $BSp(n)_2^\wedge$  is (weakly)  $N$ -determined. So  $BL_p^\wedge$  and  $BSp(n)_p^\wedge$  are homotopy equivalent.  $\square$

## REFERENCES

- [1] J. Aguadé, *Constructing modular classifying spaces*, Israel J. Math. **66** (1989), 23–40.
- [2] K. Andersen, J. Grodal, J.M. Møller, A. Viruel, *The classification of  $p$ -compact groups,  $p$  odd*, Preprint <http://arxiv.org/abs/math.AT/0302346>.
- [3] K.S. Brown, *Cohomology of groups* Springer-Verlag New York Berlin Heidelberg London Paris Tokyo Hong Kong Barcelona Budapest (1994).
- [4] A. Bousfield, D. Kan, *Homotopy limits, completion and localisation*, SLNM **304**, Springer Verlag (1972).
- [5] A. Clark, J. R. Ewing, *The realization of polynomial algebras as cohomology rings*, Pacific J. Math. **50** (1974), 425–434.

- [6] M. Curtis, A. Wiederhold, B. Williams, *Normalizers of maximal tori*, Lecture Notes in Math. **418**, 31–47.
- [7] , W.G. Dwyer, *Lie groups and  $p$ -compact groups*, Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998). Doc. Math. 1998, Extra Vol. II, 433–442 (electronic).
- [8] W.G. Dwyer, C.W. Wilkerson, *The center of a  $p$ -compact group*, The Čech centennial (Boston, MA, 1993), 119–157, Contemp. Math., 181, Amer. Math. Soc., Providence, RI, 1995.
- [9] W.G. Dwyer, C.W. Wilkerson, *Homotopy fixed point methods for Lie groups and finite loop space*, Ann. Math. **139** (1994), 395–442.
- [10] W.G. Dwyer, C.W. Wilkerson,  *$p$ -compact groups with abelian Weyl groups*, Preprint.
- [11] W.G. Dwyer, C.W. Wilkerson, *A new finite loop space at the prime two*, J. Amer. Math. Soc. **6** (1993), 37–63.
- [12] W.G. Dwyer, C.W. Wilkerson, *Product splitting for  $p$ -compact groups*, Fund. Math. **147** (1995), no. 3, 279–300.
- [13] W.G. Dwyer, C.W. Wilkerson, *A cohomology decomposition theorem*, Topology **31** (1992), 433–443.
- [14] W.G. Dwyer, A. Zabrodsky, *Maps between classifying spaces*, Algebraic topology, Barcelona, 1986, 106–119, Lecture Notes in Math., 1298, Springer, Berlin, 1987.
- [15] J.H. Gunawardena, J. Lannes, S. Zarati, *Cohomology des groupes symétriques et application de Quillen*, Advances in homotopy theory (Cortona, 1988) 61–68, London Math. Soc. Lecture Note Ser., 139.
- [16] S. Jackowski, J. McClure, R. Oliver, *Homotopy classification of self-maps of  $BG$  via  $G$ -actions, part I and part II*, Ann. Math. **135** (1992), 183–270.
- [17] J. Lannes, *Sur les espaces fonctionnelles dont la source est la classifiant d'un  $p$ -groupe abélien élémentaire* Publ. Math. IHES **75** (1992), 135–244.
- [18] J.M. Møller, *Normalizers of maximal tori* Math. Z. **231** (1999), no. 1, 51–74.
- [19] J.M. Møller, *Rational isomorphism of  $p$ -compact groups*, Topology **35** (1996), no. 1, 201–225.
- [20] J.M. Møller, *Deterministic  $p$ -compact groups*, Stable and unstable homotopy (Toronto, ON, 1996), 255–278, Fields Inst. Commun., 19, Amer. Math. Soc., Providence, RI, 1998.
- [21] J.M. Møller, *Homotopy Lie groups*, Bull. Amer. Math. Soc. **32** (1995), no. 4, 413–428.
- [22] J.M. Møller, D. Notbohm, *Centers and finite coverings of finite loop spaces*, J. Reine Angew. Math. **456** (1994), 99–133.
- [23] J.M. Møller, D. Notbohm, *Connected finite loop space with maximal tori*, Trans. Amer. Math. Soc. **350** (1998), no. 9, 3483–3504.
- [24] D. Notbohm, *Homotopy uniqueness of classifying spaces of compact connected Lie groups at primes dividing the order of the Weyl group*, Topology **33** (1994), no. 2, 271–330.
- [25] D. Notbohm, *A uniqueness result for orthogonal groups as 2-compact groups*, Arch. Math. (Basel) **78** (2002), no. 2, 110–119.
- [26] D. Notbohm, *On the 2-compact group  $DI(4)$* , Preprint.
- [27] D. Notbohm, *Spaces with polynomial mod- $p$  cohomology*, Proc. Cambridge Philos. Soc. **126** (1999), 277–292.
- [28] D. Notbohm, *Classifying spaces of compact Lie groups and finite loop spaces*, Handbook of Algebraic Topology, North-Holland (1995).
- [29] D. Notbohm, *Maps between classifying spaces*, Math. Z. **207** (1991), 229–257.
- [30] D. Notbohm, *Fake Lie groups with maximal tori IV*, Math. Ann. **294** (1992), no. 1, 109–116.
- [31] R. Oliver,  *$p$ -stubborn groups of the classical compact Lie groups*, Journal of Pure and Applied Algebra **92**, 55–78.
- [32] D. Quillen, *On the cohomology and  $K$ -theory of the general linear group over a finite field*, Ann. Math. **96** (1972), 552–586.



- [33] A. Vavpetič, A. Viruel, *On the homotopy type of the classifying space of the exceptional Lie group of rank 4*, Manuscripta Math. **107** (2002), no. 4, 521–540.
- [34] A. Viruel, *Homotopy uniqueness of  $BG_2$* , Manuscripta Math. **95** (1998), no. 4, 471–497.
- [35] C.W. Wilkerson, *Rational maximal tori*, J. Pure Appl. Algebra **4** (1974), 261–272.
- [36] Z. Wojtkowiak, *On maps from  $\mathrm{holim} F$  to  $Z$* , in Algebraic Topology, Barcelona 1986, SLNM **1298**, 227–236.
- [37] A. Zabrodsky, *On the realization of invariant subgroups of  $\pi_*(X)$* , Trans. Amer. Math. Soc. **285** (1984), 467–496.

FAKULTETA ZA MATEMATIKO IN FIZIKO, UNIVERZA V LJUBLJANA, JADRANSKA 19, SI-1111 LJUBLJANA, SLOVENIA

*E-mail address:* ales.vavpetic@FMF.Uni-Lj.Si

DPTO DE ÁLGEBRA, GEOMETRÍA Y TOPOLOGÍA, UNIVERSIDAD DE MÁLAGA, APDO CORREOS 59, 29080 MÁLAGA, SPAIN

*E-mail address:* viruel@agt.cie.uma.es